Author Index

Abu-Zeid, M. M., see Yoshioka, M. 133
Anraku, Y., see Snada, S. 223
Arai, Y.
——, Hirukawa, M. and Hanai, T.
Retention of aromatic acids on an octadecyl-bonded vinyl alcohol copolymer gel 27
——, Yamaguchi, J. and Hanai, T.
Enthalpy effect in the retention of aromatic acids on an octadecyl-bonded silica gel 21
Araki, M., see Tanaka, N. 33
Asibal, C. F., see Holzer, G. 317
Atamna, I., see Grushka, E. 5
Bayer, E.
—— and Paulus, A.
Silanophilic interactions in reversed-phase high-performance liquid chromatography 1
Brown, P. R., see Simpson, R. C. 297
Byun, D.-S., see Kasama, T. 241
Dean, T. A., see Poole, S. K. 323
Egeler, E., see Vogt, W. 83
Eisenbeiss, F., see Vogt, W. 83
Evershed, R. P.
——, Male, V. L. and Goad, L. J.
Strategy for the analysis of steryl esters from plant and animal tissues 187
Fujimori, H., see Yoshioka, M. 133
Fuke, A., see Yoshioka, M. 133
Fukui, T., see Makino, K. 271
Goad, L. J., see Evershed, R. P. 187
Görög, S.
—— and Herényi, B.
Analysis of steroids. XXXVIII. The use of high-performance liquid chromatography with diode-array UV detection for estimating impurity profiles of steroid drugs 177
Goto, A., see Yoshioka, M. 133
Grushka, E.
—— and Atamna, I.
Isochronal optimization. III. Experimental verification 5
Haginaka, J.
——, Wakai, J., Yasuda, H., Uno, T., Takahashi, K. and Katagi, T.
High-performance liquid chromatographic determination of ampicillin and its metabolites in rat plasma, bile and urine by post-column degradation with sodium hypochlorite 101
Hanai, T., see Arai, Y. 21, 27
Hara, S., see Ohkuma, T. 47
Harada, K.-I., see Oka, H. 253
Hasegawa, M., see Yuki, Y. 65
Hashimoto, Y., see Moriyasu, M. 143
Hashizume, K., see Tanaka, N. 33
Hayashi, A., see Sasaki, H. 123
Herényi, B., see Görög, S. 177
Hirai, K., see Yoshioka, M. 133
Hirukawa, M., see Arai, Y. 27
Holzer, G.
——, Zalkow, L. H. and Asibal, C. F.
Capillary supercritical fluid chromatography of pyrrolizidine alkaloids 317
Hoshino, T.
——, Suzuki, M., Ysukawa, K. and Takeuchi, M.
Effects of detergent on the elution profiles of latex beads in sedimentation field-flow fractionation 361
Ikai, Y., see Oka, H. 253
Ikawa, S.
——, Miyake, M., Mura, T. and Ikeguchi, M.
High-performance liquid chromatographic–fluorescence determination of human faecal bile acids 149
Ikeguchi, M., see Ikawa, S. 149
Imai, K.
——, Matsunaga, Y., Tsukamoto, Y. and Nishitani, A.
Application of bis[4-nitro-2-(3,6,9-trioxadecyloxycarbonyl)phenyl] oxalate to post-column chemiluminescence detection in high-performance liquid chromatography 169
Imaishi, H., see Makino, K. 271
Inoue, N., see Takeuchi, M. 207
Iriyama, K.
——, Iwamoto, T. and Yoshiura, M.
Electrochemically treated glassy carbon electrode for amperometric detection in high-performance liquid chromatography 263
Ishii, M., see Yoshioka, M. 133
Ito, N.
——, Noguchi, K., Kazama, M. and Kasai, K.-I.
Analysis of human glutamyl- and lysylplasminogen by high-performance affinity chromatography 163
Iwamori, M., see Snada, S. 223
Iwamoto, T., see Iriyama, K. 263
Izawa, A., see Snada, S., 223
Kalinoski, H. T.
——, Udseth, H. R., Wright, B. W. and Smith, R. D.
Analytical applications of capillary supercritical fluid chromatography–mass spectrometry 307

Kasai, K.-I., see Ito, N. 163
——, see Shimura, K. 353
Kasama, T.
——, Byun, D.-S. and Seyama, Y.
Quantitative analysis of sterols in serum by high-performance liquid chromatography. Application to the biochemical diagnosis of cerebrotendinous xanthomatosis 241
Katagi, T., see Haginaka, J. 101
Kato, A., see Moriyasu, M. 143
Kawamura, N., see Oka, H. 253
Kazama, M., see Ito, N. 163
Kim, K. R.
——, Lee, Y. J., Lee, H. S. and Zlatkis, A.
Solid-phase extraction of chloramphenicol with graphitized carbon black 285
Kimoto, H., see Yuki, Y. 65
Kinoshita, T., see Kobayashi, Y. 113
Kitagaki-Ogawa, H., see Matsumoto, I. 77
——, see Sasaki, H. 123
Kobata, A., see Takeuchi, M. 207
Kobayashi, E., see Suzuki, K. T. 233
Kobayashi, Y.
——, Kubo, H. and Kinoshita, T.
Post-column derivatization system for the fluorimetric determination of guanidino compounds with ninhydrin by reversed-phase ion-pair high-performance liquid chromatography 113
Kosuge, Y., see Terada, H. 343
Koyama, T., see Matsumoto, I. 77
Kubo, H., see Kobayashi, Y. 113
Lee, H. S., see Kim, K. R. 285
Lee, Y. J., see Kim, K. R. 285
Makino, K.
——, Ozaki, H., Matsumoto, T., Imaishi, H., Takeuchi, T. and Fukui, T.
Reversed-phase ion-pair chromatography of oligodeoxyribonucleotides 271
Male, V. L., see Evershed, R. P. 187
Matsumoto, I.
——, Koyama, T., Kitagaki-Ogawa, H. and Seno, N.
Separation is isolectins by high-performance hydrophobic interaction chromatography 77
——, see Sasaki, H. 123
Matsumoto, T., see Makino, K. 271
Matsunaga, Y., see Imai, K. 169
Meyer, H. D., see Vogt, W. 83
Miyake, M., see Ikawa, S. 149
Mohri, Z.-I., see Yagi, S. 279
Moriyasu, M.
——, Kato, A. and Hashimoto, Y.
Kinetic studies of fast equilibrium by means of high-performance liquid chromatography. XV. Separation of tautomers of β-diketoesters 143
Motamura, H., see Sakuma, S. 293
Mura, T., see Ikawa, S. 149
Murayama, W., see Terada, H. 343
Nagai, Y., see Snada, S. 223
Nakagawa, T., see Yagi, S. 297
Nakaya, N., see Terada, H. 343
Nambara, T., see Shimada, K. 215
Nishitani, A., see Imai, K. 169
Niwa, T., see Shimada, K. 215
Noguchi, K., see Ito, N. 163
Nunogaki, K.-I., see Terada, H. 343
Nunogaki, Y., see Terada, H. 343
Ohkuma, T.
—— and Hara, S.
Tail-producing slow adsorption–desorption process in liquid–solid chromatography 47
Oka, H.
——, Ikai, Y., Kawamura, N., Uno, K., Yamada, M., Harada, K.-I and Suzuki, M.
Improvement of chemical analysis of antibiotics. XII. Simultaneous analysis of seven tetracyclines in honey 253
Ozaki, H., see Makino, K. 271
Parvez, H., see Yoshioka, M. 133
Paulus, A., see Bayer, E. 1
Poole, C. F., see Poole, S. K. 323
Poole, S. K.
——, Dean, T. A. and Poole, C. F.
Preparation of environmental samples for the determination of polycyclic aromatic hydrocarbons by thin-layer chromatography 323
Qureshi, G. A.
High-performance liquid chromatographic methods with fluorescence detection for the determination of branched-chain amino acids and their alpha-keto analogues in plasma samples of healthy subjects and uraemic patients 91
—— and P. Södersten
Determination of free amino acids in the cerebrospinal fluid or male rats by high-performance liquid chromatography with fluorescence detection. Alterations in amino acid concentrations during sexual activity 247
Saigo, K., see Yuki, Y. 65
Sakuma, S.
—— and Motomura, H.
Purification of saikosaponins a, c and d. Application of large-scale reversed-phase high-performance liquid chromatography 293

Sasaki, H.
——, Hayashi, A., Kitagaki-Ogawa, H., Matsumoto, I. and Seno, N.
Improved method for the immobilization of heparin 123
Seno, N., see Matsumoto, I. 77
——, see Sasaki, H. 123
Seyama, Y., see Kasama, T. 241
Shimada, K.
——, Xie, F., Niwa, T., Wakasawa, T. and Nambara, T.
Studies on steroids. CCXXIX. Separation and characterization of catechol oestrogen glucuronides in urine of pregnant women by high-performance liquid chromatography 215
Shimura, K.
—— and Kasai, K.-I.
Affinophoresis of pea lectin and fava bean lectin with an anionic affinophore, bearing *p*-aminophenyl-α-D-mannoside as an affinity ligand 353
Simpson, R. C.
—— and Brown, P. R.
Fast, microbore, and fast microbore high-performance liquid chromatography of nucleic acid constituents 297
Smith, R. D., see Kalinoski, H. T. 307
Snada, S.
——, Uchida, Y., Anraku, Y., Izawa, A., Iwamori, M. and Nagai, Y.
Analysis of ceramide and monohexaosyl glycolipid derivatives by high-performance liquid chromatography and its application to the determination of the molecular species in tissues 223
Södersten, P., see Qureshi, G. A. 247
Sommer, W., see Vogt, W. 83
Sugihira, N., see Suzuki, K. T. 233
Sugimoto, T., see Yoshioka, M. 133
Sunaga, H., see Suzuki, K. T. 233
Suzuki, K. T.
——, Sunaga, H., Kobayashi, E. and Sugihira, N.
High-performance liquid chromatography–inductively coupled plasma profiles of cadmium, zinc, sulphur and other elements in rat liver supernatants after cadmium injection 233
Suzuki, M., see Hoshino, T. 361
——, see Oka, H. 253
Tachibana, K., see Yuki, Y. 65
Takahashi, K., see Haginaka, J. 101
Takasaki, S., see Takeuchi, M. 207
Takeuchi, M.
——, Takasaki, S., Inoue, N. and Kobata, A.
Sensitive method for carbohydrate composition analysis of glycoproteins by high-performance liquid chromatography 207
——, see Hoshino, T. 361
Takeuchi, T., see Makino, K. 271
Tanaka, H., see Yagi, S. 279
Tanaka, N.
——, Hashizume, K. and Araki, M.
Comparison of polymer-based stationary phases with silica-based stationary phases in reversed-phase liquid chromatography. Selective binding of rigid, compact molecules by alkylated polymer gels 33
Terada, H.
——, Kosuge, Y., Murayama, W., Nakaya, N., Nunogaki, Y. and Nunogaki, K.-I.
Correlation of hydrophobic parameters of organic compounds determined by centrifugal partition chromatography with partition coefficients between octanol and water 343
Tsukamoto, Y., see Imai, K. 169
Uchida, Y., see Snada, S. 223
Udseth, H. R., see Kalinoski, H. T. 307
Uno, K., see Oka, H. 253
Uno, T., see Haginaka, J. 101
Vogt, W.
——, Egeler, E., Sommer, W., Eisenbeiss, F. and Meyer, H. D.
High-performance liquid chromatographic determination of hormonal peptides and their fluorenylmethoxycarbonyl derivatives 83
Wakai, J., see Haginaka, J. 101
Wakasawa, T., see Shimada, K. 215
Wright, B. W., see Kalinoski, H. T. 307
Xie, F., see Shimada, K. 215
Yagi, S.
——, Nakagawa, T., Tanaka, H., Yoshitake, A. and Mohri, Z.-I.
High-performance liquid chromatographic system for the rapid purification of growth-hormone receptor in rat livers 297
Yamada, K., see Yoshioka, M. 133
Yamada, M., see Oka. H. 253
Yamaguchi, J., see Arai, Y. 21
Yasuda, H., see Haginaka, J. 101
Yoshioka, M.
——, Yamada, K., Abu-Zeid, M. M., Fujimori, H., Fuke, A., Hirai, K., Goto, A., Ishii, M., Sugimoto, T. and Parvez, H.
Anlyses of adenosine and adenine nucleotides in biological materials by fluorescence reaction–high-performance liquid chromatography 133
Yoshitake, A., see Yagi, S. 279
Yoshiura, M., see Iriyama, K. 263

Ysukawa, K., see Hoshino, T. 361
Yuki, Y.
——, Saigo, K., Kimoto, H.,Tachibana, K. and Hasegawa, M.
Novel chiral stationary phases for optical resolution by ligand-exchange high-performance liquid chromatography 65

Zalkow, L. H., see Holzer, G. 317
Zlatkis, A., see Kim, K. R. 285